

Choiceless Set-theoretic Geology

Toshimichi Usuba (薄葉季路)

Waseda University (早稲田大学)

June. 21, 2019

Asian Logic Conference 2019

Nazarbayev University, Nursultan, Kazakhstan

Abstract

Set-theoretic geology is a study of the structure of all ground models of the universe V . We will try to extend the standard set-theoretic geology to choiceless set-theoretic geology, which can treat with choiceless models.

Definability of ground models

Throughout this talk, forcing means a **set forcing**.

Fact (Laver, Woodin)

In any forcing extension $V[G]$ of V , the universe V is a (first order) definable class in $V[G]$ with some parameters from V .

Changing the viewpoint

Every ground model of V is definable by some first order formula of set-theory.

A **ground** is a ground model of V .

Definability of ground models

Throughout this talk, forcing means a [set forcing](#).

Fact (Laver, Woodin)

In any forcing extension $V[G]$ of V , the universe V is a (first order) definable class in $V[G]$ with some parameters from V .

Changing the viewpoint

Every ground model of V is definable by some first order formula of set-theory.

A [ground](#) is a ground model of V .

Definability of ground models

Throughout this talk, forcing means a [set forcing](#).

Fact (Laver, Woodin)

In any forcing extension $V[G]$ of V , the universe V is a (first order) definable class in $V[G]$ with some parameters from V .

Changing the viewpoint

Every ground model of V is definable by some first order formula of set-theory.

A [ground](#) is a ground model of V .

Uniform definability of grounds

Actually all grounds can be defined by some **uniform** way.

Fact (Fuchs-Hamkins-Reitz)

There is a first order formula $\varphi(x, y)$ of set theory such that:

- 1 *For each set r , the class $W_r = \{x \mid \varphi(x, r)\}$ is a ground of V with $r \in W_r$ ($W_r = V$ is possible).*
- 2 *For every transitive model $M \subseteq V$ of ZFC, if M is a ground of V , then there is r with $M = W_r$.*

Remark

The formula φ does not depend on V : In any model M of ZFC, φ defines its grounds by the same way.

Set-theoretic geology

- - ▶ The intersection of two grounds.
 - ▶ Whether every ground satisfies some statement?
 - ▶ Whether every ground has a proper ground?
 - ▶ The intersection of all grounds.
- These are second-order objects and second-order statements of set-theory.
- However, the uniform definability allows us to describe such statements in ZFC, and study the structure of the collection of grounds $\{W_r \mid r \in V\}$ in ZFC: e.g.,
 - ▶ One can define (in ZFC) the intersection of two grounds.
 - ▶ One can ask (in ZFC) if every ground satisfies some statement.
 - ▶ One can ask (in ZFC) whether $\forall r \exists s (W_s \subsetneq W_r)$.
 - ▶ One can define (in ZFC) the intersection of all grounds, and ask (in ZFC) whether what is that?

This study is now called **set-theoretic geology**.

Set-theoretic geology

- - ▶ The intersection of two grounds.
 - ▶ Whether every ground satisfies some statement?
 - ▶ Whether every ground has a proper ground?
 - ▶ The intersection of all grounds.
- These are second-order objects and second-order statements of set-theory.
- However, the uniform definability allows us to describe such statements in ZFC, and study the structure of the collection of grounds $\{W_r \mid r \in V\}$ in ZFC: e.g.,
 - ▶ One can define (in ZFC) the intersection of two grounds.
 - ▶ One can ask (in ZFC) if every ground satisfies some statement.
 - ▶ One can ask (in ZFC) whether $\forall r \exists s (W_s \subsetneq W_r)$.
 - ▶ One can define (in ZFC) the intersection of all grounds, and ask (in ZFC) whether what is that?

This study is now called **set-theoretic geology**.

Geological facts: Mantle

An important geological object is the mantle:

Definition

The **mantle** \mathbb{M} is the intersection of all grounds.

\mathbb{M} is a parameter free definable transitive class containing all ordinals. The mantle is the **core** of all grounds.

Fact (Fuchs-Hamkins-Reitz, U.)

The mantle is a model of ZFC, and is the largest forcing invariant definable model;

- *For every ground W and forcing extension $V[G]$, we have $\mathbb{M}^W = \mathbb{M}^{V[G]} = \mathbb{M}$,*
- *If N is a definable transitive model of ZFC satisfying $N^W = N^{V[G]} = N$ for every W and $V[G]$, then $N \subseteq \mathbb{M}$.*

Geological facts: Mantle

An important geological object is the mantle:

Definition

The **mantle** \mathbb{M} is the intersection of all grounds.

\mathbb{M} is a parameter free definable transitive class containing all ordinals. The mantle is the **core** of all grounds.

Fact (Fuchs-Hamkins-Reitz, U.)

The mantle is a model of ZFC, and is the largest forcing invariant definable model;

- *For every ground W and forcing extension $V[G]$, we have $\mathbb{M}^W = \mathbb{M}^{V[G]} = \mathbb{M}$,*
- *If N is a definable transitive model of ZFC satisfying $N^W = N^{V[G]} = N$ for every W and $V[G]$, then $N \subseteq \mathbb{M}$.*

Geological facts: Mantle

An important geological object is the mantle:

Definition

The **mantle** \mathbb{M} is the intersection of all grounds.

\mathbb{M} is a parameter free definable transitive class containing all ordinals. The mantle is the **core** of all grounds.

Fact (Fuchs-Hamkins-Reitz, U.)

The mantle is a model of ZFC, and is the largest forcing invariant definable model;

- *For every ground W and forcing extension $V[G]$, we have $\mathbb{M}^W = \mathbb{M}^{V[G]} = \mathbb{M}$,*
- *If N is a definable transitive model of ZFC satisfying $N^W = N^{V[G]} = N$ for every W and $V[G]$, then $N \subseteq \mathbb{M}$.*

Geological facts: DDG

A key was the **downward directedness of grounds**.

Fact (U.)

*If M and N are grounds of V , then M and N have a common ground.
More strongly, for every family of set-many grounds $\{W_r \mid r \in X\}$, there is W which is a ground of each W_r ($r \in X$).*

DDG yields:

- For every α , the α -initial segment of the mantle \mathbb{M}_α is of the form W_α for some ground W .
- Consequently, the mantle is a model of ZFC.
- The mantle is forcing invariant.

Geological facts: DDG

A key was the **downward directedness of grounds**.

Fact (U.)

*If M and N are grounds of V , then M and N have a common ground.
More strongly, for every family of set-many grounds $\{W_r \mid r \in X\}$, there is W which is a ground of each W_r ($r \in X$).*

DDG yields:

- For every α , the α -initial segment of the mantle \mathbb{M}_α is of the form W_α for some ground W .
- Consequently, the mantle is a model of ZFC.
- The mantle is forcing invariant.

Geological facts: Manipulating Mantle 1

Fact (Fuchs-Hamkins-Reitz)

The universe can be the mantle of some class forcing extension: There is a class forcing extension $V[G]$ of V such that $V = \mathbb{M}^{V[G]}$.

Hence, unlike L and other canonical inner models, $\text{Th}(\mathbb{M})$ cannot be determined.

Geological facts: Manipulating Mantle 2

\mathbb{M} can be far from, or close to V as you like.

Fact (Fuchs-Hamkins-Reitz)

- 1 V has a class forcing extension $V[G]$ such that $V[G]$ has no proper ground, so $V[G] = \mathbb{M}^{V[G]}$.
- 2 V has a class forcing extension $V[G]$ such that $V[G]$ is not a forcing extension of its mantle.

However Large cardinal gives some restriction:

Fact (U.)

If there exists a very large cardinal (extendible cardinal), then the mantle is the minimum ground of V , hence V is a forcing extension of the mantle.

Geological facts: Manipulating Mantle 2

\mathbb{M} can be far from, or close to V as you like.

Fact (Fuchs-Hamkins-Reitz)

- 1 V has a class forcing extension $V[G]$ such that $V[G]$ has no proper ground, so $V[G] = \mathbb{M}^{V[G]}$.
- 2 V has a class forcing extension $V[G]$ such that $V[G]$ is not a forcing extension of its mantle.

However Large cardinal gives some restriction:

Fact (U.)

If there exists a very large cardinal (extendible cardinal), then the mantle is the minimum ground of V , hence V is a forcing extension of the mantle.

Set-theoretic geology without AC

- In the standard geology, the universe V and all grounds are supposed to satisfy the Axiom of Choice (AC).
- However it is possible that V has a ground which does not satisfy AC.
- Moreover choiceless context is very common (e.g., $ZF+AD$), and forcing over choiceless model is very useful (e.g., \mathbb{P}_{max} over $L(\mathbb{R})$).
- So it is natural to extend the standard set-theoretic geology to **Choiceless Set-theoretic Geology**, which can treat with such choiceless models.

Question

Can we develop choiceless set-theoretic geology?

- A first problem is the definability of grounds of V .

Set-theoretic geology without AC

- In the standard geology, the universe V and all grounds are supposed to satisfy the Axiom of Choice (AC).
- However it is possible that V has a ground which does not satisfy AC.
- Moreover choiceless context is very common (e.g., $ZF+AD$), and forcing over choiceless model is very useful (e.g., \mathbb{P}_{max} over $L(\mathbb{R})$).
- So it is natural to extend the standard set-theoretic geology to **Choiceless Set-theoretic Geology**, which can treat with such choiceless models.

Question

Can we develop choiceless set-theoretic geology?

- A first problem is the definability of grounds of V .

Set-theoretic geology without AC

- In the standard geology, the universe V and all grounds are supposed to satisfy the Axiom of Choice (AC).
- However it is possible that V has a ground which does not satisfy AC.
- Moreover choiceless context is very common (e.g., ZF+AD), and forcing over choiceless model is very useful (e.g., \mathbb{P}_{max} over $L(\mathbb{R})$).
- So it is natural to extend the standard set-theoretic geology to **Choiceless Set-theoretic Geology**, which can treat with such choiceless models.

Question

Can we develop choiceless set-theoretic geology?

- A first problem is the definability of grounds of V .

Set-theoretic geology without AC

- In the standard geology, the universe V and all grounds are supposed to satisfy the Axiom of Choice (AC).
- However it is possible that V has a ground which does not satisfy AC.
- Moreover choiceless context is very common (e.g., $ZF+AD$), and forcing over choiceless model is very useful (e.g., \mathbb{P}_{max} over $L(\mathbb{R})$).
- So it is natural to extend the standard set-theoretic geology to **Choiceless Set-theoretic Geology**, which can treat with such choiceless models.

Question

Can we develop choiceless set-theoretic geology?

- A first problem is the definability of grounds of V .

Set-theoretic geology without AC

- In the standard geology, the universe V and all grounds are supposed to satisfy the Axiom of Choice (AC).
- However it is possible that V has a ground which does not satisfy AC.
- Moreover choiceless context is very common (e.g., $ZF+AD$), and forcing over choiceless model is very useful (e.g., \mathbb{P}_{max} over $L(\mathbb{R})$).
- So it is natural to extend the standard set-theoretic geology to **Choiceless Set-theoretic Geology**, which can treat with such choiceless models.

Question

Can we develop choiceless set-theoretic geology?

- A first problem is the definability of grounds of V .

Set-theoretic geology without AC

- In the standard geology, the universe V and all grounds are supposed to satisfy the Axiom of Choice (AC).
- However it is possible that V has a ground which does not satisfy AC.
- Moreover choiceless context is very common (e.g., $ZF+AD$), and forcing over choiceless model is very useful (e.g., \mathbb{P}_{max} over $L(\mathbb{R})$).
- So it is natural to extend the standard set-theoretic geology to **Choiceless Set-theoretic Geology**, which can treat with such choiceless models.

Question

Can we develop choiceless set-theoretic geology?

- A first problem is the definability of grounds of V .

Covering and approximation properties

In ZFC, Hamkins' covering and approximation properties are important tools for proving the (uniform) definability of grounds.

Definition (Hamkins)

Let κ be a cardinal, and $M \subseteq V$ a transitive (set or class) model of ZFC.

- 1 M satisfies **the κ -covering property** if for every $a \in [\text{ON}]^{<\kappa}$ there exists $b \in M \cap [\text{ON}]^{<\kappa}$ with $a \subseteq b$.
- 2 M satisfies **the κ -approximation property** if for every set A of ordinals, $\forall x \in [\text{ON}]^{<\kappa} \cap M (A \cap x \in M)$ then $A \in M$.

Fact (Hamkins)

Let M, N be transitive models of ZFC. If M and N satisfy the κ -covering and the κ -approximation properties, and $\mathcal{P}(\kappa^+) \cap M = \mathcal{P}(\kappa^+) \cap N$, then $M \cap \mathcal{P}(\text{ON}) = N \cap \mathcal{P}(\text{ON})$, hence $M = N$.

Covering and approximation properties

In ZFC, Hamkins' covering and approximation properties are important tools for proving the (uniform) definability of grounds.

Definition (Hamkins)

Let κ be a cardinal, and $M \subseteq V$ a transitive (set or class) model of ZFC.

- 1 M satisfies **the κ -covering property** if for every $a \in [\text{ON}]^{<\kappa}$ there exists $b \in M \cap [\text{ON}]^{<\kappa}$ with $a \subseteq b$.
- 2 M satisfies **the κ -approximation property** if for every set A of ordinals, $\forall x \in [\text{ON}]^{<\kappa} \cap M (A \cap x \in M)$ then $A \in M$.

Fact (Hamkins)

Let M, N be transitive models of ZFC. If M and N satisfy the κ -covering and the κ -approximation properties, and $\mathcal{P}(\kappa^+) \cap M = \mathcal{P}(\kappa^+) \cap N$, then $M \cap \mathcal{P}(\text{ON}) = N \cap \mathcal{P}(\text{ON})$, hence $M = N$.

Fact (Hamkins)

Every ground satisfies the κ -covering and approximation properties for some κ .

Each ground W can be defined as: For a large κ , let $X = \mathcal{P}(\kappa^+) \cap W$. Then W is definable with the parameter $r = \langle \kappa, X, \mathbb{P}, G \rangle$ as the unique model M satisfying the κ -covering, κ -approximation, $M \cap \mathcal{P}(\kappa^+) = X$, and $V = M[G]$.

- In the proof of the uniform definability of grounds, the property:

$$M \cap \mathcal{P}(ON) = N \cap \mathcal{P}(ON) \Rightarrow M = N$$

is essential. However this is not valid if AC fails in M and N .

- Moreover, it is not clear that, in ZF, every ground satisfies the covering and approximation properties.

Fact (Hamkins)

Every ground satisfies the κ -covering and approximation properties for some κ .

Each ground W can be defined as: For a large κ , let $X = \mathcal{P}(\kappa^+) \cap W$. Then W is definable with the parameter $r = \langle \kappa, X, \mathbb{P}, G \rangle$ as the unique model M satisfying the κ -covering, κ -approximation, $M \cap \mathcal{P}(\kappa^+) = X$, and $V = M[G]$.

- In the proof of the uniform definability of grounds, the property:

$$M \cap \mathcal{P}(ON) = N \cap \mathcal{P}(ON) \Rightarrow M = N$$

is essential. However this is not valid if AC fails in M and N .

- Moreover, it is not clear that, in ZF, every ground satisfies the covering and approximation properties.

Fact (Hamkins)

Every ground satisfies the κ -covering and approximation properties for some κ .

Each ground W can be defined as: For a large κ , let $X = \mathcal{P}(\kappa^+) \cap W$. Then W is definable with the parameter $r = \langle \kappa, X, \mathbb{P}, G \rangle$ as the unique model M satisfying the κ -covering, κ -approximation, $M \cap \mathcal{P}(\kappa^+) = X$, and $V = M[G]$.

- In the proof of the uniform definability of grounds, the property:

$$M \cap \mathcal{P}(ON) = N \cap \mathcal{P}(ON) \Rightarrow M = N$$

is essential. However this is not valid if AC fails in M and N .

- Moreover, it is not clear that, in ZF, every ground satisfies the covering and approximation properties.

Question (in ZF, open)

- 1 For every poset \mathbb{P} and generic G , is V definable in $V[G]$?
- 2 Are all (choiceless) grounds of V uniformly definable?

Fact (Gitman-Johnstone, in ZF)

Suppose DC_κ holds. Then for every poset \mathbb{P} with size $\leq \kappa$ (\mathbb{P} is assumed to be well-orderable), V is definable in $V^{\mathbb{P}}$ with some parameters.

Their result does not need the full AC but a weak AC, and it is not known if \mathbb{P} is not well-orderable.

Question (in ZF, open)

- 1 For every poset \mathbb{P} and generic G , is V definable in $V[G]$?
- 2 Are all (choiceless) grounds of V uniformly definable?

Fact (Gitman-Johnstone, in ZF)

Suppose DC_κ holds. Then for every poset \mathbb{P} with size $\leq \kappa$ (\mathbb{P} is assumed to be well-orderable), V is definable in $V^{\mathbb{P}}$ with some parameters.

Their result does not need the full AC but a weak AC, and it is not known if \mathbb{P} is not well-orderable.

Convention

- From now on, our base theory is ZF unless otherwise specified.
- A **ground** is a transitive model of ZF such that V is a set-generic extension of it.

Löwenheim-Skolem property in ZF

For the uniform definability of grounds, we give a partial answer: It is possible if some Löwenheim-Skolem property holds.

Definition

We say that κ is a **Löwenheim-Skolem cardinal** if for every $\alpha > \kappa$, $p \in V_\alpha$, and $\gamma < \kappa$, there is $\beta > \alpha$ and $X \prec V_\beta$ such that:

- 1 $p \in X$ and $V_\gamma \subseteq X$.
- 2 $V_\gamma X \subseteq X$.
- 3 The transitive collapse of X belongs to V_κ .

Remark

- In ZFC, κ is Löwenheim-Skolem if and only if $\beth_\kappa = \kappa$, so there are always proper class many Löwenheim-Skolem cardinals.
- If κ is a limit of Löwenheim-Skolem cardinals, then κ is Löwenheim-Skolem as well.

Löwenheim-Skolem property in ZF

For the uniform definability of grounds, we give a partial answer: It is possible if some Löwenheim-Skolem property holds.

Definition

We say that κ is a **Löwenheim-Skolem cardinal** if for every $\alpha > \kappa$, $p \in V_\alpha$, and $\gamma < \kappa$, there is $\beta > \alpha$ and $X \prec V_\beta$ such that:

- 1 $p \in X$ and $V_\gamma \subseteq X$.
- 2 $V_\gamma X \subseteq X$.
- 3 The transitive collapse of X belongs to V_κ .

Remark

- In ZFC, κ is Löwenheim-Skolem if and only if $\beth_\kappa = \kappa$, so there are always proper class many Löwenheim-Skolem cardinals.
- If κ is a limit of Löwenheim-Skolem cardinals, then κ is Löwenheim-Skolem as well.

Löwenheim-Skolem property in ZF

For the uniform definability of grounds, we give a partial answer: It is possible if some Löwenheim-Skolem property holds.

Definition

We say that κ is a **Löwenheim-Skolem cardinal** if for every $\alpha > \kappa$, $p \in V_\alpha$, and $\gamma < \kappa$, there is $\beta > \alpha$ and $X \prec V_\beta$ such that:

- 1 $p \in X$ and $V_\gamma \subseteq X$.
- 2 $V_\gamma X \subseteq X$.
- 3 The transitive collapse of X belongs to V_κ .

Remark

- In ZFC, κ is Löwenheim-Skolem if and only if $\beth_\kappa = \kappa$, so there are always proper class many Löwenheim-Skolem cardinals.
- If κ is a limit of Löwenheim-Skolem cardinals, then κ is Löwenheim-Skolem as well.

Some consequences of LS property

The existence of a Löwenheim-Skolem cardinal is not provable from ZF:

Lemma

If κ is a singular Löwenheim-Skolem cardinal, (e.g., singular limit of Löwenheim-Skolem cardinals), then κ^+ is regular, and the non-stationary ideal over κ^+ is κ^+ -complete. In addition for every regressive function $f : \kappa^+ \rightarrow \kappa^+$, there is $\gamma < \kappa^+$ such that the set $\{\alpha < \kappa^+ \mid f(\alpha) = \gamma\}$ is stationary in κ^+ .

In Gitik's model with no regular uncountable cardinals, Löwenheim-Skolem cardinal does not exist. Compare:

Fact (Woodin)

If κ is a singular limit of supercompact cardinals, then κ^+ is regular, and the non-stationary ideal over κ^+ is κ^+ -complete.

Some consequences of LS property

The existence of a Löwenheim-Skolem cardinal is not provable from ZF:

Lemma

If κ is a singular Löwenheim-Skolem cardinal, (e.g., singular limit of Löwenheim-Skolem cardinals), then κ^+ is regular, and the non-stationary ideal over κ^+ is κ^+ -complete. In addition for every regressive function $f : \kappa^+ \rightarrow \kappa^+$, there is $\gamma < \kappa^+$ such that the set $\{\alpha < \kappa^+ \mid f(\alpha) = \gamma\}$ is stationary in κ^+ .

In Gitik's model with no regular uncountable cardinals, Löwenheim-Skolem cardinal does not exist. Compare:

Fact (Woodin)

If κ is a singular limit of supercompact cardinals, then κ^+ is regular, and the non-stationary ideal over κ^+ is κ^+ -complete.

Some consequences of LS property

The existence of a Löwenheim-Skolem cardinal is not provable from ZF:

Lemma

If κ is a singular Löwenheim-Skolem cardinal, (e.g., singular limit of Löwenheim-Skolem cardinals), then κ^+ is regular, and the non-stationary ideal over κ^+ is κ^+ -complete. In addition for every regressive function $f : \kappa^+ \rightarrow \kappa^+$, there is $\gamma < \kappa^+$ such that the set $\{\alpha < \kappa^+ \mid f(\alpha) = \gamma\}$ is stationary in κ^+ .

In Gitik's model with no regular uncountable cardinals, Löwenheim-Skolem cardinal does not exist. Compare:

Fact (Woodin)

If κ is a singular limit of supercompact cardinals, then κ^+ is regular, and the non-stationary ideal over κ^+ is κ^+ -complete.

Large cardinal yields LS property

Definition (Woodin)

An uncountable cardinal κ is **supercompact** if for every $\alpha > \kappa$, there is $\beta > \alpha$, a transitive set X , and an elementary embedding $j : V_\beta \rightarrow X$ such that the critical point of j is κ , $\alpha < j(\kappa)$, and $V_\alpha X \subseteq X$.

Lemma

Every supercompact cardinal is Löwenheim-Skolem, and a limit of Löwenheim-Skolem cardinals.

Hence if there are proper class many supercompact cardinals, then there are proper class many Löwenheim-Skolem cardinals as well.

Large cardinal yields LS property

Definition (Woodin)

An uncountable cardinal κ is **supercompact** if for every $\alpha > \kappa$, there is $\beta > \alpha$, a transitive set X , and an elementary embedding $j : V_\beta \rightarrow X$ such that the critical point of j is κ , $\alpha < j(\kappa)$, and $V_\alpha X \subseteq X$.

Lemma

Every supercompact cardinal is Löwenheim-Skolem, and a limit of Löwenheim-Skolem cardinals.

Hence if there are proper class many supercompact cardinals, then there are proper class many Löwenheim-Skolem cardinals as well.

Uniform definability of grounds

Under the assumption that there are proper class many Löwenheim-Skolem cardinals, we can define all grounds uniformly.

Theorem

Suppose there are proper class many Löwenheim-Skolem cardinals. Then all grounds are uniformly definable: There is a first order formula $\varphi(x, y)$ of set-theory such that:

- 1 For each set r , the class $W_r = \{x \mid \varphi(x, r)\}$ is a ground of V with $r \in W_r$.
- 2 For every ground $M \subseteq V$, there is r with $M = W_r$.

Under reasonable assumption, we can start the [choiceless set-theoretic geology](#).

Uniform definability of grounds

Under the assumption that there are proper class many Löwenheim-Skolem cardinals, we can define all grounds uniformly.

Theorem

Suppose there are proper class many Löwenheim-Skolem cardinals. Then all grounds are uniformly definable: There is a first order formula $\varphi(x, y)$ of set-theory such that:

- 1 For each set r , the class $W_r = \{x \mid \varphi(x, r)\}$ is a ground of V with $r \in W_r$.
- 2 For every ground $M \subseteq V$, there is r with $M = W_r$.

Under reasonable assumption, we can start the [choiceless set-theoretic geology](#).

Coarse measure

Let us outline the proof of the uniform definability of grounds. For the sake of this, we introduce alternative measure on sets, which is coarser than the standard cardinality but works without AC.

Definition

For a set x , let $\|x\|$, the **norm** of x , be the least ordinal α such that there is a surjection from V_α onto x .

Remark

- 1 $\|x\| \leq \text{rank}(x)$.
- 2 If $x \subseteq y$ then $\|x\| \leq \|y\|$.
- 3 If $M \subseteq V$ is a transitive model of (a sufficiently large fragment of) ZF and $x \in M$, then $\|x\| \leq \|x\|^M$.
- 4 κ is Löwenheim-Skolem \iff every first order structure has an elementary substructure with some closure property and norm $< \kappa$.

Coarse measure

Let us outline the proof of the uniform definability of grounds. For the sake of this, we introduce alternative measure on sets, which is coarser than the standard cardinality but works without AC.

Definition

For a set x , let $\|x\|$, the **norm** of x , be the least ordinal α such that there is a surjection from V_α onto x .

Remark

- 1 $\|x\| \leq \text{rank}(x)$.
- 2 If $x \subseteq y$ then $\|x\| \leq \|y\|$.
- 3 If $M \subseteq V$ is a transitive model of (a sufficiently large fragment of) ZF and $x \in M$, then $\|x\| \leq \|x\|^M$.
- 4 κ is Löwenheim-Skolem \iff every first order structure has an elementary substructure with some closure property and norm $< \kappa$.

Coarse measure

Let us outline the proof of the uniform definability of grounds. For the sake of this, we introduce alternative measure on sets, which is coarser than the standard cardinality but works without AC.

Definition

For a set x , let $\|x\|$, the **norm** of x , be the least ordinal α such that there is a surjection from V_α onto x .

Remark

- 1 $\|x\| \leq \text{rank}(x)$.
- 2 If $x \subseteq y$ then $\|x\| \leq \|y\|$.
- 3 If $M \subseteq V$ is a transitive model of (a sufficiently large fragment of) ZF and $x \in M$, then $\|x\| \leq \|x\|^M$.
- 4 κ is Löwenheim-Skolem \iff every first order structure has an elementary substructure with some closure property and norm $< \kappa$.

Covering and approximation properties

Definition

Let M be a transitive (set or class) model of (a sufficiently large fragment of) ZF, and α an ordinal with $\alpha \in M$.

- 1 M satisfies the α -norm covering property if for every $\beta \in M$ and $x \subseteq M_\beta$, if $\|x\| < \alpha$ then there is $y \in M$ such that $x \subseteq y$ and $\|y\|^M < \alpha$.
- 2 M satisfies the α -norm approximation property if for every $\beta \in M$ and $A \subseteq M_\beta$, if $A \cap x \in M$ for every $x \in M$ with $\|x\|^M < \alpha$, then $A \in M$.

Uniqueness of models with covering and approximation properties

Theorem

Let κ be a Löwenheim-Skolem cardinal, and M and N be transitive models of (a sufficiently large fragment of) ZF. If

- 1 $\kappa \in M \cap N$ and $M \cap ON = N \cap ON$.
- 2 M and N satisfy the γ -norm covering and approximation properties for some $\gamma < \kappa$.
- 3 $M_\kappa = N_\kappa$.

Then $M = N$.

By induction on the rank of sets.

Lemma

Suppose κ is a Löwenheim-Skolem cardinal. For every ground M of V , if there is a poset $\mathbb{P} \in M_\kappa$ such that V is a generic extension of M via \mathbb{P} , then M satisfies the κ -norm covering and approximation properties.

Theorem (Uniform definability of all grounds)

Suppose there are proper class many Löwenheim-Skolem cardinals. Then there is a first order formula $\varphi(x, y)$ of set theory such that:

- 1 For each set r , the class $W_r = \{x \mid \varphi(x, r)\}$ is a ground of V with $r \in W_r$ ($W_r = V$ is possible).
- 2 For every transitive model $M \subseteq V$ of ZFC, if M is a ground of V , then there is r with $M = W_r$.

Hence all grounds are uniformly definable.

Lemma

Suppose κ is a Löwenheim-Skolem cardinal. For every ground M of V , if there is a poset $\mathbb{P} \in M_\kappa$ such that V is a generic extension of M via \mathbb{P} , then M satisfies the κ -norm covering and approximation properties.

Theorem (Uniform definability of all grounds)

Suppose there are proper class many Löwenheim-Skolem cardinals. Then there is a first order formula $\varphi(x, y)$ of set theory such that:

- 1 For each set r , the class $W_r = \{x \mid \varphi(x, r)\}$ is a ground of V with $r \in W_r$ ($W_r = V$ is possible).
- 2 For every transitive model $M \subseteq V$ of ZFC, if M is a ground of V , then there is r with $M = W_r$.

Hence all grounds are uniformly definable.

Generic absoluteness of LS property

Lemma

Suppose $M \subseteq V$ is a ground, κ a cardinal, and $V = M[G]$ for some $G \subseteq \mathbb{P} \in M_\kappa$.

- 1 If κ is a limit of Löwenheim-Skolem cardinals, then κ is a Löwenheim-Skolem cardinal in M .
- 2 In M , if κ is a limit of Löwenheim-Skolem cardinals, then κ is a Löwenheim-Skolem cardinal in V .

Corollary

The statement “there are proper class many Löwenheim-Skolem cardinals” is absolute between V , all grounds, and all generic extensions.

Generic absoluteness of LS property

Lemma

Suppose $M \subseteq V$ is a ground, κ a cardinal, and $V = M[G]$ for some $G \subseteq \mathbb{P} \in M_\kappa$.

- 1 If κ is a limit of Löwenheim-Skolem cardinals, then κ is a Löwenheim-Skolem cardinal in M .
- 2 In M , if κ is a limit of Löwenheim-Skolem cardinals, then κ is a Löwenheim-Skolem cardinal in V .

Corollary

The statement “there are proper class many Löwenheim-Skolem cardinals” is absolute between V , all grounds, and all generic extensions.

Mantle

As in ZFC, we can define the mantle of V .

Definition

The **mantle** \mathbb{M} is the intersection of all grounds (which may or may not satisfy AC).

Question

Is the mantle a model of ZF (or ZFC)?

In ZFC, key property was the downward directedness of grounds (DDG). However the downward directedness of grounds can fail in ZF.

Lemma

It is consistent that V satisfies AC, but V has two grounds M and N such that AC fails in M and N , and M and N have no common ground. In addition the mantle of M (and N) is a model of ZFC.

Mantle

As in ZFC, we can define the mantle of V .

Definition

The **mantle** \mathbb{M} is the intersection of all grounds (which may or may not satisfy AC).

Question

Is the mantle a model of ZF (or ZFC)?

In ZFC, key property was the downward directedness of grounds (DDG). However the downward directedness of grounds can fail in ZF.

Lemma

It is consistent that V satisfies AC, but V has two grounds M and N such that AC fails in M and N , and M and N have no common ground. In addition the mantle of M (and N) is a model of ZFC.

Mantle

As in ZFC, we can define the mantle of V .

Definition

The **mantle** \mathbb{M} is the intersection of all grounds (which may or may not satisfy AC).

Question

Is the mantle a model of ZF (or ZFC)?

In ZFC, key property was the downward directedness of grounds (DDG). However the downward directedness of grounds can fail in ZF.

Lemma

It is consistent that V satisfies AC, but V has two grounds M and N such that AC fails in M and N , and M and N have no common ground. In addition the mantle of M (and N) is a model of ZFC.

If AC is forceable

In ZFC, there are proper class many Löwenheim-Skolem cardinals.

Corollary

If there is a poset which forces AC (AC is *forceable*), then there are proper class many Löwenheim-Skolem cardinals, and all grounds are uniformly definable.

Hence a natural question arises: Under what circumstances is AC forceable?

Fact (Blass)

The following are equivalent:

- 1 *AC is forceable.*
- 2 *There is a set X such that for every set Y , there is an ordinal α and a surjection from $X \times \alpha$ onto Y .*

If AC is forceable

In ZFC, there are proper class many Löwenheim-Skolem cardinals.

Corollary

If there is a poset which forces AC (AC is **forceable**), then there are proper class many Löwenheim-Skolem cardinals, and all grounds are uniformly definable.

Hence a natural question arises: Under what circumstances is AC forceable?

Fact (Blass)

The following are equivalent:

- 1 *AC is forceable.*
- 2 *There is a set X such that for every set Y , there is an ordinal α and a surjection from $X \times \alpha$ onto Y .*

If AC is forceable

In ZFC, there are proper class many Löwenheim-Skolem cardinals.

Corollary

If there is a poset which forces AC (AC is **forceable**), then there are proper class many Löwenheim-Skolem cardinals, and all grounds are uniformly definable.

Hence a natural question arises: Under what circumstances is AC forceable?

Fact (Blass)

The following are equivalent:

- ① *AC is forceable.*
- ② *There is a set X such that for every set Y , there is an ordinal α and a surjection from $X \times \alpha$ onto Y .*

If AC is forceable

In ZFC, there are proper class many Löwenheim-Skolem cardinals.

Corollary

If there is a poset which forces AC (AC is **forceable**), then there are proper class many Löwenheim-Skolem cardinals, and all grounds are uniformly definable.

Hence a natural question arises: Under what circumstances is AC forceable?

Fact (Blass)

The following are equivalent:

- 1 *AC is forceable.*
- 2 *There is a set X such that for every set Y , there is an ordinal α and a surjection from $X \times \alpha$ onto Y .*

Another characterization

For a transitive model $M \subseteq V$ of ZF and set X , let $M(X)$ be the minimal transitive model of ZF containing $M \cup \{X\}$.

Theorem

The following are equivalent:

- 1 There is a poset which forces AC.
- 2 There is a definable transitive model W of ZFC and a set X such that $W(X) = V$.
- 3 There is a definable transitive model W of ZFC and a set X such that $W(X) = V$, and W is a ground of some generic extension of V .

AC is forceable $\iff V$ is a **small** extension of a model of ZFC.

Another characterization

For a transitive model $M \subseteq V$ of ZF and set X , let $M(X)$ be the minimal transitive model of ZF containing $M \cup \{X\}$.

Theorem

The following are equivalent:

- 1 There is a poset which forces AC.
- 2 There is a definable transitive model W of ZFC and a set X such that $W(X) = V$.
- 3 There is a definable transitive model W of ZFC and a set X such that $W(X) = V$, and W is a ground of some generic extension of V .

AC is forceable $\iff V$ is a **small** extension of a model of ZFC.

Geology under AC forceable

Corollary

Suppose AC is forceable.

- 1 Every ground of V is of the form $M(X)$ for some definable transitive model M of ZFC and set X , and M is a ground of some generic extension of V (M may not be a ground of V).
- 2 For every two grounds M and N , there is $W \subseteq M \cap N$ which is a ground of some generic extension of V (but W may not be a ground of V , M , and N).

Corollary

Suppose V satisfies AC, and $W \subseteq V$ is a transitive model of ZF. Then the following are equivalent:

- 1 W is a ground of V .
- 2 There is a ground M of V such that M satisfies AC and $W = M(X)$ for some $X \in M$.

In particular,

- The family of grounds satisfying AC is dense in all grounds, with respects to \subseteq .
- The mantle (the intersection of all grounds which may or may not satisfy AC) is a model of ZFC.
- If V is a generic extension of the mantle \mathbb{M} , then for every $M \subseteq V$, M is ground of V if and only if M is of the form $\mathbb{M}(X)$ for some X .

Questions: Definability of grounds

Question

In ZF, are all grounds always uniformly definable?

If not, AC cannot be forceable over the universe. Some known candidates:

- A model with no uncountable regular cardinals (Gitik).
- A model which has proper class many infinite but Dedekind-finite sets (Monro).
- A model in which Fodor's lemma fails everywhere, and every non-stationary ideal is not σ -complete (Karagila).
- ...
- Bristol model M , which is a transitive model of ZF with $L \subseteq M \subseteq L^{\mathbb{C}}$, definable in $L^{\mathbb{C}}$, but $M \neq L(X)$ for every set X . AC is not forceable over M .

Questions: Geology of specific models

- A model with no uncountable regular cardinals (Gitik).
- A model which has a proper class many infinite but Dedekind-finite sets (Monro).
- A model in which Fodor's lemma fails everywhere, and every non-stationary ideal is not σ -complete (Karagila).
- Bristol model M , which is a model of ZF with $L \subseteq M \subseteq L^{\mathbb{C}}$, but $M \neq L(X)$ for every set X . AC is not forceable over M .

How is the geology of such models?

Test questions

- 1 Are grounds of these models uniformly definable? Or Do these models have proper class many Löwenheim-Skolem cardinals?
- 2 Does these models have a proper ground?
- 3 What is the mantle of these models?

Questions: AC-conjecture

If AC is forceable, then the geology of V is trivial: it is almost the same to the geology with AC. So we want to know non-trivial situation, but the failure of AC is sensitive under our setting.

Theorem (Woodin, in essence)

- 1 If there is a Löwenheim-Skolem cardinal, then DC is forceable.
- 2 If there are proper class many regular Löwenheim-Skolem cardinals, then there is a **class forcing** which forces AC.

Question

Is it consistent that there are proper class many Löwenheim-Skolem cardinals, but AC is never forceable?

This question may be connected with Woodin's AC-Conjecture.

Conjecture (Woodin)

If there are large cardinals, then AC is forceable.

Questions: AC-conjecture

If AC is forceable, then the geology of V is trivial: it is almost the same to the geology with AC. So we want to know non-trivial situation, but the failure of AC is sensitive under our setting.

Theorem (Woodin, in essence)

- 1 If there is a Löwenheim-Skolem cardinal, then DC is forceable.
- 2 If there are proper class many regular Löwenheim-Skolem cardinals, then there is a **class forcing** which forces AC.

Question

Is it consistent that there are proper class many Löwenheim-Skolem cardinals, but AC is never forceable?

This question may be connected with Woodin's AC-Conjecture.

Conjecture (Woodin)

If there are large cardinals, then AC is forceable.

Questions: AC-conjecture

If AC is forceable, then the geology of V is trivial: it is almost the same to the geology with AC. So we want to know non-trivial situation, but the failure of AC is sensitive under our setting.

Theorem (Woodin, in essence)

- 1 If there is a Löwenheim-Skolem cardinal, then DC is forceable.
- 2 If there are proper class many regular Löwenheim-Skolem cardinals, then there is a **class forcing** which forces AC.

Question

Is it consistent that there are proper class many Löwenheim-Skolem cardinals, but AC is never forceable?

This question may be connected with Woodin's AC-Conjecture.

Conjecture (Woodin)

If there are large cardinals, then AC is forceable.

Questions: Geological

There are many geological questions.

Question

- 1 Is the mantle a model of ZF? (or ZFC?)
- 2 Is it consistent that the mantle is a model ZF but not ZFC?

Fact (Fuchs-Hamkins-Reitz, in ZFC)

The universe can be the mantle of some class forcing extension: There is a class forcing extension $V[G]$ of V such that $V = \mathbb{M}^{V[G]}$.

Question

Does ZF-version of Fuchs-Hamkins-Reitz's theorem hold?

Questions: Geological

There are many geological questions.

Question

- 1 Is the mantle a model of ZF? (or ZFC?)
- 2 Is it consistent that the mantle is a model ZF but not ZFC?

Fact (Fuchs-Hamkins-Reitz, in ZFC)

The universe can be the mantle of some class forcing extension: There is a class forcing extension $V[G]$ of V such that $V = \mathbb{M}^{V[G]}$.

Question

Does ZF-version of Fuchs-Hamkins-Reitz's theorem hold?

Questions: Geological

Fact (Fuchs-Hamkins-Reitz, in ZFC)

- 1 V has a class forcing extension $V[G]$ such that $V[G]$ has no proper ground, so $V[G] = \mathbb{M}^{V[G]}$.
- 2 It is consistent that V is not a forcing extension of \mathbb{M} .

Fact (U., in ZFC)

If there exists an extendible cardinal, then the mantle is the minimum ground of V , hence V is a forcing extension of the mantle.

Question

In ZF, do similar results hold?

Questions: DDG

DDG can fail in choiceless models, however, in many cases, for every two ground M and N we can find a model W such that $W \subseteq M \cap N$ and W is a **generic ground**, a ground of some generic extension of V .

Question

Does the following weak form of DDG hold?

For every grounds M and N of V , there is a generic ground W with $W \subseteq M \cap N$.

- If AC is forceable then this weak DDG holds.
- On the other hand, even if this weak DDG holds, it is not clear that the mantle is a model of ZF.

References

- A. Blass, *Injectivity, projectivity, and the axiom of choice*. Trans. Amer. Math. Soc. 255 (1979), 31–59.
- G. Fuchs, J. D. Hamkins, J. Reitz, *Set-theoretic geology*. Ann. Pure Appl. Logic 166 (2015), no. 4, 464–501.
- M. Gitik, *All uncountable cardinals can be singular*. Israel J. Math. 35 (1980), no. 1-2, 61–88.
- S. Grigorieff, *Intermediate submodels and generic extensions in set theory*. Annal. Math. Vol. 101 (1975), No. 3, 447–490 .
- V. Gitman, T. Johnstone, *On ground model definability*. Infinity, computability, and metamathematics, 205–227, Tributes, 23, Coll. Publ., London, 2014.
- J. D. Hamkins, *Extensions with the approximation and cover properties have no new large cardinals*. Fund. Math. 180 (2003), no. 3, 257–277.
- A. Karagila, *Fodor's lemma can fail everywhere*. Acta Math. Hungarica, Vol. 154 (2018), Issue 1, 231–242.

References

- A. Karagila, *The Bristol Model: an abyss called a Cohen real*. Journal of Mathematical Logic, Vol. 18 (2018), No. 02, 1850008.
- R. Laver, *Certain very large cardinals are not created in small forcing extensions*. Ann. Pure Appl. Logic 149 (2007), no. 1-3, 1–6.
- G. P. Monro, *Independence results concerning Dedekind-finite sets*. J. Austral. Math. Soc. 19 (1975), 35–46.
- J. Reitz, *The Ground Axiom*. J. of Symbolic Logic 72 (2007), no. 4, 1299–1317.
- T. Usuba, *The downward directed grounds hypothesis and very large cardinals*. J. Math. Log. 17 (2017), no. 2, 1750009, 24 pp.
- T. Usuba, *Choiceless Löwenheim-Skolem property and uniform definability of grounds*. <https://arxiv.org/abs/1904.00895>
- W. H. Woodin, *Suitable extender models I*. J. Math. Log. 10 (2010), no. 1-2, 101–339.

Thank you for your attention!